arXiv:astro-ph/0302597v2 10 Mar 2003

WMAP data and the curvature of space

Jean-Philippe Uzan
Laboratoire de Physique Théorique, CNRS–UMR 8627,
Bât. 210, Université Paris XI, F-91405 Orsay Cedex, France,
Institut d'Astrophysique de Paris, GReCO, CNRS-FRE 2435, 98 bis, Bd Arago, 75014 Paris, France,
Department of Mathematics and Applied Mathematics,
University of Capetown, Rondebosch 7700, Capetown, South Africa

Ulrich Kirchner and George F.R. Ellis
Department of Mathematics and Applied Mathematics,
University of Capetown, Rondebosch 7700, Capetown, South Africa

(Dated: February 28, 2003)

Inter alia, the high precision WMAP data on Cosmic Background Radiation marginally indicate that the universe has positively curved (and hence spherical) spatial sections. In this paper we take this data seriously and consider some of the consequences for the background dynamics. In particular, we show that this implies a limit to the number of e-foldings that could have taken place in the inflationary epoch; however this limit is consistent with some inflationary models that solve all the usual cosmological problems and are consistent with standard structure formation theory.

The Wilkinson Microwave Anisotropy Probe (WMAP) has recently provided high resolution Cosmic Microwave Background data [1, 2] of importance for cosmology - indeed they constitute a significant contribution towards the goal of developing precision cosmology. Among the interesting conclusions that have been reached from this data are constraints on present value Ω_0 of the total density parameter of the universe. The WMAP results interestingly indicate that while the universe is close to being flat ($\Omega_0 \simeq 1$), a closed universe is marginally preferred: $\Omega_0 > 1$ [1]. In particular, with a prior on the Hubble constant, one gets that $\Omega_0 = 1.03 \pm 0.05$ at 95% confidence level, while combining WMAP data with type Ia supernovae leads to $\Omega_0 = 1.02 \pm 0.04$ or to $\Omega_0 = 1.02 \pm 0.02$ respectively without and with a prior on the Hubble parameter. The latter may be regarded as the present best estimate of this parameter. Note also that this tendency to point toward a closed universe is not strictly speaking either new or restricted to the WMAP data. For instance with a prior on the nature of the initial conditions, the Hubble parameter and the age of the universe, analysis of the DASI, BOOMERanG, MAXIMA and DMR data [3, 4] lead to $\Omega_0 = 0.99 \pm 0.12$ but to $\Omega_0 = 1.04 \pm 0.05$ if one takes into account only the DASI, BOOMERanG and CBI data, both at 1σ-level. The Archeops balloon experiment [5] points toward $\Omega_0 = 1.00^{+0.03}_{-0.02}$ with data from the HST and a prior on the Hubble constant but to values in the range $\Omega_0 = 1.16^{+0.24}_{-0.20}$ from combined CBR data alone and to $\Omega_0 = 1.04^{+0.02}_{-0.04}$ from combined CBR and supernova data, all at a 68% confidence level. The improved precision from WMAP is clear.

The confirmation of the existence of Doppler peaks and their respective locations tends to confirm the inflationary paradigm [6], as does the existence of an almost scale invariant power spectrum. Furthermore a nearly flat universe, argued to be predicted generically by most inflationary models [7], is usually seen as a further evidence. Since

$$\delta(t) \equiv \Omega(t) - 1 = \frac{K}{a^2 H^2} \tag{1}$$

where $a(t)$ is the scale function and $H(t) = \dot{a}/a$ the Hubble parameter, the WMAP constraints on the curvature of space imply that the curvature density parameter, $\Omega_K(t) \equiv K/a^2H^2$, has been smaller than a few percent from the time the largest observable scale, $k_{\min} \sim a_0 H_0$, crossed the horizon during inflation up to today. It is then a good approximation to neglect the curvature from the time $k_{\min}$ crossed the horizon to today when performing structure formation calculations, mainly because $\Omega_{k_{\min}} = \Omega_{\text{today}}$ (see Fig. 1). The constraints on the curvature of the universe thus back up the validity of calculations underlying the tests of inflation performed up to now that assume a flat universe and focus on the properties of the perturbations (spectral indices, tensor modes, statistics, etc.).

Nevertheless, when $K \neq 0$ (which is highly favoured over the case $K = 0$ by probability considerations: the latter are of measure zero in the space of Friedman-Lemaître-Robertson-Walker universes in most measures), Ω_K is driven toward 0 exponentially during inflation. The curvature term therefore was necessarily dominant at early enough times, if inflation undergoes a sufficient number of e-foldings, mainly because the curvature term behaves as a^{-2} and will tend to dominate over any kind of matter having an equation of state less stiff than $-1/3$, and thus over a slow rolling scalar field or a cosmological constant, in the early inflationary era.

We argue that the study of dynamics of the background for such models can give interesting insight into the inflationary era and its dynamics prior to $k_{\min}$ horizon exit. Working backward in time from the present, we aim to emphasize the constraints that arise on the inflationary models from the WMAP data. In doing so, we take seriously the indication that a value of Ω_0 slightly larger than 1 is favoured. During a slow-roll inflationary phase, the universe is described by an almost de Sitter spacetime with the $K=+1$ scale factor

$$a(t) \propto \cosh(Ht) \tag{2}$$

instead of the exponential growth obtained in the flat case. It follows that the number of e-foldings between the onset and the end of inflation cannot be arbitrarily large, when we take present day cosmological parameters into account. The curvature enhances the curvature, $\ddot{a}$, of the scale factor in the early stage of inflation so that there exists a turn-around point in the past. It follows [8, 9] that the number of e-foldings during inflation is bounded by a maximum possible number of e-foldings, $N_{\rm max}$, related to the present day density parameter Ω_0.

Assuming that the inflation is driven simply by a cosmological constant, the scale factor of a closed universe in this epoch is given by

$$a(t) = a_{\rm min}\cosh(\lambda t) \tag{3}$$

with $\lambda \equiv \sqrt{\Lambda/3}$ and $t=0$ corresponds to the minimum expansion of the universe. It follows that $\delta = \Omega - 1 = 1/\sinh^2(\lambda {\rm t})$. The time t_* at which $k_{\rm min} = a_0 H_0$ crosses the horizon is thus obtained to be

$$\lambda t_* = {\rm Arcsinh}(1/\sqrt{\delta_0}). \tag{4}$$

The number of e-foldings between $t=0$ and t_* is simply

$$\Delta N_{\rm max} = \ln\frac{a_{k_{\rm min}}}{a_{\rm min}} \tag{5}$$

that is

$$\Delta N_{\rm max} = \frac{1}{2}\ln\left(1+\frac{1}{\delta_0}\right). \tag{6}$$

It typically ranges from 2.3 to 1.5 when δ_0 varies from 0.01 to 0.05 (see Fig. 2).

This argument also holds for negatively curved universes, in which case, the scale factor behaves as $a \propto \sinh\lambda t$. In that case, there will not be any maximal number of e-foldings but the universe starts off from a singularity. This will enhance the trans-Planckian problem [10].

Another way to look at the existence of a maximum number of e-foldings in the $K=+1$ case, is to relate the number of e-foldings during inflation to the later history of the universe [8, 9]. During inflation, the comoving Hubble length aH is decreasing, while it is increasing at all further times. A given comoving scale, k, crosses the horizon when $k = a(t_k)H(t_k) \equiv a_k H_k$. The number of e-foldings, $N(k)$, between when this scale crosses the horizon and the end of inflation is obtained to be

$$N(k) = \ln\left(\frac{a_{\rm end}H_{\rm end}}{a_k H_k}\right). \tag{7}$$

The determination of $N(k)$ requires that we compute

$$\mathcal{R} \equiv \frac{a_0}{a_{\rm end}} \tag{8}$$

which requires a complete history of the evolution of the universe from the end of inflation to today. If one assumes that the universe is matter dominated from the end of (slow roll) inflation to reheating, radiation dominated from reheating to equality and matter dominated up to now [27], then [11, 12]

$$\begin{aligned} N(k) \;&\simeq\; 62 - \ln\left(\frac{k}{a_0H_0}\right) - \ln\left(\frac{10^{16}\,{\rm GeV}}{V_k^{1/4}}\right) \\ &+\frac{1}{4}\ln\frac{V_k}{V_{\rm end}} - \frac{1}{3}\ln\left(\frac{V_{\rm end}^{1/4}}{\rho_{\rm reh}^{1/4}}\right) - \ln h, \end{aligned} \tag{9}$$

h being the Hubble parameter in unites of 100 km.s^{-1}/Mpc. For our purpose, it will be more convenient to relate $N(k)$ to $\mathcal{R}$ which can easily be found to be given by

$$-\ln\mathcal{R} \simeq -66 - \frac{1}{3}\ln\left(\frac{V_{\rm end}^{1/4}}{\rho_{\rm reh}^{1/4}}\right) + \ln\left(\frac{10^{16}\,{\rm GeV}}{V_{\rm end}^{1/4}}\right). \tag{10}$$

It follows that

$$\begin{aligned} N(k) \;&\simeq\; 128 - \ln\mathcal{R} - \ln\left(\frac{k}{a_0H_0}\right) \\ &-2\ln\left(\frac{10^{16}\,{\rm GeV}}{V_k^{1/4}}\right) - \ln h. \end{aligned} \tag{11}$$

The cosmic microwave background roughly probe scales from 10 Mpc (i.e. of the order of the thickness of the last scattering surface) to 10^4 Mpc (i.e. of the order of the size of the observable universe) while galaxy surveys can probe the range 1 Mpc to 100 Mpc.

Concerning the characteristic scales involved in the problem, we can assume that $V_k \sim V_{\rm end}$ as long as slow rolling holds. The reheating temperature can be argued to be larger than $\rho_{\rm reh}^{1/4} > 10^{10}$ GeV to avoid the gravitino problem [13] and may be pushed in the extreme case to 10^3 GeV, i.e. just before the electro-weak transition so, that baryogenesis can take place. The amplitude of the cosmological fluctuations (typically of order 2×10^{-5} on Hubble scales) roughly implies that $V_{\rm end}^{1/4}$ is smaller than a few times 10^{16} GeV and, for the same reason as above, has to be larger than 10^{10} GeV in the extreme case.

This implies that the number of e-foldings has approximately to lie between 50 and 70, which is the order of magnitude also required to solve the horizon and flatness problem when $K=0$ [7], but it can be lowered to 25 in the extreme case of thermal inflation [14], which we are not considering here.

In the following, we restrict our analysis to the case where reheating takes place just at the end of inflation, i.e. $\rho_{\rm reh} = V_{\rm end}$. It has been shown [8] that, in that case, the existence of a positive curvature limits the number of e-foldings to

$$N_{\rm max}(\mathcal{R},\delta_0)=\ln\sqrt{\frac{\alpha}{\delta_0}} \tag{12}$$

with

$$\alpha=(1+\delta_0-\Omega_{\rm rad0})\mathcal{R}+\Omega_{\rm rad0}\mathcal{R}^2 \tag{13}$$

where $\delta_0\equiv\Omega_0-1$ and $\Omega_{\rm rad0}\simeq 4.17\times10^{-5}h^{-2}$ is the radiation density parameter today. Let us emphasize that the derivation of Eq. (12), as well as Eq. (6), does not make any assumption concerning what happened before the start of inflation.

This allows us to estimate the number of e-foldings allowed prior to the time at which $k_{\rm min}$ crossed the horizon,

$$\Delta N=N_{\rm max}[\mathcal{R}(V_{\rm end}^{1/4}),\delta_0]-N(k_{\rm min},V_{\rm end}^{1/4}) \tag{14}$$

where we will let δ_0 vary in the range $0.01-0.05$ (see Fig. 2). It can be checked, as expected from Eq. (12) that ΔN depends slightly on $V_{\rm end}^{1/4}$.

We see that the allowed number of e-foldings are compatible with the requirements of structure formation. 'Flatness' is of course solved to the accuracy represented by δ_0. The scenarios sketched here do not lead to as small a value of δ_0 as is often supposed - and thereby is compatible with the best-fit WMAP data. However, for the mean value of the curvature, one obtains that

$$\delta_0=0.02\Longleftrightarrow\Delta N_{\rm max}=1.97. \tag{15}$$

With standard parameter values this gives the maximum possible number of e-foldings as about $N_{\rm max}=62+1.97\simeq64$ - compatible with the estimates given above, but not nearly as large as suggested in some models of inflation.

In view of the bounds on $N_{\rm max}$, there are also bounds on the ability of inflation to solve the horizon problem when one considers inflationary dynamics with $K=+1$ [9]. Roughly speaking, the horizon problem is only solved during the inflationary era itself if the start of the inflationary era is close enough to the throat at $t=0$ in the expansion (see Eq. 3); that is, if the total number N of e-foldings during inflation is close to the total $N_{\rm max}$ allowed, i.e. $N\simeq N_{\rm max}$. If inflation starts sufficiently far from that throat, then it does not matter how many e-foldings take place, they cannot provide causal connectivity on large enough scale to solve the horizon problem; rather that connectivity has to be set up in the pre-inflationary era (and is evidenced by a smooth enough small patch that is then expanded to large scales by inflation). However inflation certainly can start near enough the throat to solve the horizon problem during the inflationary era and still lead to $\delta_0=0.02$. Examples of non-slow rolling cases are the emergent and awakening universes [15] that start form an Einstein static state and are singularity free. The above formula (15) limiting the number of e-foldins will not apply in these cases.



FIG. 1: The curvature $(1/a^2H^2)$ as a function of time (schematically). The solid line represents the case of a flat universe ($K=0$)with exponential inflation while the dashed line depicts the case of a closed universe ($K=+1$). Before the end of inflation aH scales either as $e^{\lambda t}$ ($K=0$) or $1/\sinh\lambda t$ ($K=+1$) while it scales as $\alpha t^{\alpha-1}$ during matter and radiation dominated eras. The largest observed scale is depicted as well, and one can easily see that Ω_K will be smaller than a few percent between the time it crosses the horizon during inflation to today when it reenters the horizon.



FIG. 2: The maximum number of e-foldings allowed before the largest observable scale, $k_{\rm min}$ exits the horizon during inflation, as a function of δ_0.

In this note we have argued that the curvature of space may play an important role in the dynamics of inflation in its early stage. We have argued that a maximum number of $1.5-2.5$ e-foldings can take place before the largest scale observed in the universe crossed the horizon during slow-roll inflation. However, despite the fact that the curvature term will be smaller than a few percent during all times ranging from the horizon exit of the largest observable mode to its reentry today, a positive curvature term as small as the one indicated by current data will dramatically change the early phase of inflation and hence lead to the limits on the number of e-foldings allowed as described above. These restrict the families of inflationary universes compatible with the current data, but do not exclude inflationary models even though the value of δ_0 indicated by WMAP is larger than supposed in most inflationary scenarios.

Note a possible way around the argument. In the case that inflation is driven by a scalar field, the existence of a maximal number of e-foldings may be avoided if it exits the slow-roll regime. It was however shown that the dynamics of closed universes filled with a scalar field is chaotic [16, 17, 18] and can lead to the existence of singularity free universes [19] with periodic [20] or aperiodic [16] trajectories. In that case, again, a positive curvature affects the dynamics of the early phase of inflation and the slow-roll regime has to be left before $N_{\max}$ is reached, which is also at odds with the standard picture.

Let us finally consider further features of spherical universes that may turn to out be useful in understanding the WMAP data. Surprisingly, the WMAP angular correlation function seems to lack signal strength on angular scales larger than 60 degrees [1, 2]. This may indicate a possible discreteness of the initial power spectrum, as expected e.g. from non-standard topology. The case for topology becomes most interesting in the case of spherical spaces [21]. First, the spectrum of the Laplacian in spherical spaces is always discrete [22, 23]. Second, and as emphasized in Ref. [24], a non trivial topology is most likely to be detectable in the case of spherical spaces and examples of the resulting observational effects are discussed in Ref. [25].

The same conclusion cannot be reached for negatively curved universe. It is unlikely that we could detect a compact hyperbolic space because, for an observer at a generic point, the topology scale is comparable to the curvature radius or longer. The possibility of a detection would require the observer to sit near a short closed geodesic [26].

In conclusion, if the indication that Ω_0 differs from unity at a level of a few percent is confirmed then it may imply important results concerning the dynamics of the early stage of our universe. The $K = +1$ background de Sitter model (3) differs fundamentally from the scale-free de Sitter model in the $K = 0$ frame - not least in terms of being geodesically complete. The WMAP observation suggest the former may be the appropriate model to use in investigating early inflationary dynamics.

Acknowledgements: JPU thanks the Department of Mathematics and Applied Mathematics of the University of Cape Town for hospitality. UK acknowledges financial support from the University of Cape Town. We thank Roy Maartens for his comments.

---

[1] C.L. Bennett *et al.*, [`arXiv:astro-ph/0302207`].

[2] D.N. Spergel *et al.*, [`arXiv:astro-ph/0302209`].

[3] J. L. Sievers *et al.*, [`arXiv:astro-ph/0205387`]

[4] C.B. Netterfield *et al.*, Astrophys. J. **571**, 604 (2002).

[5] A. Benoît *et al.*, [`arXiv:astro-ph/0210306`]; A. Benoît *et al.*, [`arXiv:astro-ph/0210305`].

[6] H.V. Peiris *et al.*, [`arXiv:astro-ph/0302225`]; V. Barger, H.S. Lee, and D. Marfatia, [`arXiv:hep-ph/0302150`].

[7] A.H. Guth, Phys. Rev. D **23**, 347 (1981); A.D. Linde, Phys. Lett. B **108**, 389 (1982); S.W. Hawking, Phys. Lett. B **115**, 295 (1982); A.A. Starobinski, Phys. Lett. B **117**, 175 (1982); A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).

[8] G.F.R. Ellis *et al.*, Gen. Rel. Grav. **34**, 1445 (2002).

[9] G.F.R. Ellis *et al.*, Gen. Rel. Grav. **34**, 1461 (2002).

[10] J. Martin and R. Brandenberger, Phys. Rev. D **65**, 123501 (2001); R. Brandenberger and J. Martin, Mod. Phys. Lett. A **16**, 999 (2001); M . Lemoine *et al.*, Phys. Rev. D **65**, 023510 (2002).

[11] D. Lyth and A. Liddle, Phys. Rept. **231**, 1 (1993).

[12] J.E. Lidsey *et al.*, Rev. Mod. Phys. **69**, 373 (1997).

[13] S. Sarkar, Rep. Prog. Phys. **59**, 1493 (1996).

[14] D.H. Lyth and E.D. Stewart, Phys. Rev. D **53**, 1784 (1996); *ibid.*, Phys. Rev. Lett. **75**, 201 (1995).

[15] G.F.R. Ellis and R. Maartens, [`arXiv:gr-qc/0211082`].

[16] D. N. Page, Class. Quant. Grav. **1**, 417 (1984).

[17] A. V. Toporensky, [`arXiv:gr-qc/0011113`].

[18] N. Cornish and E. P. S. Shellard, Phys. Rev. Lett. **81**, 3571 (1998).

[19] L. Parker and S. A. Fulling, Phys. Rev. D **7**, 2357 (1973); A. A. Starobinsky, Sov. Astron. Lett. **4**,82 (1978).

[20] S. W. Hawking, in *Relativity, Group and Topology II*, edited by B. S. de Witt and R. Stora, (North Holland, Amsterdam, 1984).

[21] E. Gausmann *et al.*, Class. Quant. Grav. **18**, 5155 (2001).

[22] R. Lehoucq, J.-P. Uzan, and J. Weeks, Kodai Math. J. (to appear), [`arXiv:math.SP/0202072`]; R. Lehoucq *et al.*, Class. Quant. Grav. **19**, 4683 (2002).

[23] A. Riazuelo *et al.*, [`arXiv:astro-ph/0212223`].

[24] J. Weeks, R. Lehoucq, and J.-P. Uzan, Class. Quant. Grav. (to appear), [`arXiv:astro-ph/0209389`].

[25] A. Riazuelo *et al.*, “Cosmic microwave background maps of multi-connected spherical spaces” (in preparation).

[26] J. Weeks, Int. J. Mod. Phys. A (submitted); G.I. Gomero, M.J. Rebouças, and R. Tavakol, Class. Quant. Grav. **18**, 4461 (2001); G.I. Gomero, M.J. Rebouças, and R. Tavakol, Int. J. Mod. Phys. A **17**, 4261 (2002).

[27] The results will be only marginally affected by a late-time cosmological constant as indicated by the supernova data.